

AD-A243 954



PL-TR-91-2234


✓m
(2)

TIME DEPENDENCE OF HEAVY ION CONCENTRATION
IN THE RING CURRENT

Mary K. Hudson

Dartmouth College
Hanover, NH 03755

9 September 1991



Scientific Report No. 1

Approved for public release; distribution unlimited




PHILLIPS LABORATORY
AIR FORCE SYSTEMS COMMAND
HANSOM AIR FORCE BASE, MASSACHUSETTS 01731-5000


91-18059

01 1216 021

"This technical report has been reviewed and is approved for publication"


KEVIN J. KERNS, LT, USAF
Contract Manager


E. G. MULLEN
Branch Chief


WILLIAM SWIDER
Deputy Director

This document has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify GL/IMA, Hanscom AFB, MA 01731. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

This reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204 Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

| | | | | | |
|---|---|--|--|--|--|
| 1. AGENCY USE ONLY (Leave blank) | | 2. REPORT DATE 9 September 1991 | | 3. REPORT TYPE AND DATES COVERED Scientific Report No. 1 | |
| 4. TITLE AND SUBTITLE Time Dependence of Heavy Ion Concentration in the Ring Current | | | | 5. FUNDING NUMBERS PE 62101F PR 7601 TA 20 WU CC Contract F19628-87-K-0038 | |
| 6. AUTHOR(S) Mary K. Hudson | | | | | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Dartmouth College Hanover, NH 03755 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Phillips Laboratory Hanscom AFB, MA 01731-5000 Contract Manager: Lt Kevin Kerns/PHP | | | | 10. SPONSORING/MONITORING AGENCY REPORT NUMBER PL-TR-91-2234 | |
| 11. SUPPLEMENTARY NOTES | | | | | |
| 12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; Distribution unlimited | | | | 12b. DISTRIBUTION CODE | |
| 13. ABSTRACT (Maximum 200 words) The focus of this project has been a comprehensive study of processes responsible for the decay of the storm time ring current. We have examined the importance to ring current ion loss of interaction with waves in the Pc 1 (Hz) and Pc 5 (mHz) frequency ranges, since such waves are a common feature of the ring current region during recovery from magnetic storms. Pc 1 waves, identified as electromagnetic ion cyclotron (EMIC) waves, are generated by the loss cone distribution of ring current ions. We have performed both linear dispersion analysis and particle simulations which show the importance of the loss cone vs. bi-Maxwellian nature of ring current ion distributions in determining the growth rate, angle of propagation and polarization of EMIC waves. Pitch angle diffusion is found to be an important loss process for ring current H ⁺ . Compressional Pc 5s are found to contribute to ring current O ⁺ loss. Their frequency (2-5 mHz) is comparable to the bounce frequency of ring current O ⁺ . We have performed ring current test particle simulations in prescribed wave and background fields, including a dipole magnetic, convection and corotation electric field. We find that a drift-bounce resonant interaction causes ring current O ⁺ loss to the magnetopause in the energy range of a few tens of keV for moderate Kp (1.5 - 2). | | | | | |
| 14. SUBJECT TERMS ring current, electromagnetic ion cyclotron waves, Pc 5 micropulsations, pitch angle diffusion, radial diffusion | | | | 15. NUMBER OF PAGES 12 | |
| | | | | 16. PRICE CODE | |
| 17. SECURITY CLASSIFICATION OF REPORT Unclassified | 18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified | 19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified | 20. LIMITATION OF ABSTRACT SAR | | |

Figure Captions

Fig. 1. Plot of energy versus azimuthal angle (positive westward) or local time for an ion injected at 1800 LT. The solid line represents the minimum energy required for the ion to drift westward without Pc 5 waves. The dashed line represents the energy variation under the effect of the convection and corotation electric fields, without Pc 5 waves, for a 28 keV ion injected at 1800LT; the dotted line is for the same case with Pc 5 waves included.

Fig. 2a. The equatorial trajectories of 132 O^+ ions with different initial energies: 26.00 keV to 31.00 keV, in steps of 0.50 keV; equatorial pitch angles: 15° to 75° , in steps of 20° ; and L shells: $L = 6.0$ to $L = 7.0$, in steps of 0.5; the same initial gyrophase ($= \arctan(v_y/v_x)$), $v_x = v$, $v_y = v_z = 0$. Here the Pc 5 wave has ten different azimuthal wave numbers, ranging from $m = 95$ to $m = 105$, each with a random initial azimuthal phase and the same amplitude, which now is reduced by $1/\sqrt{10}$. The azimuthal wave number determines the perpendicular wavelength $\lambda_\perp = 2\pi LR_E/m$.

Fig. 2b. Numbers of ions in 2 keV bins, integrated over L, for parameters of (a).

| | |
|-----|-------------------------------------|
| For | |
| I | <input checked="" type="checkbox"/> |
| | <input type="checkbox"/> |
| | <input type="checkbox"/> |
| Ion | |

| | |
|--------------------|----------------------|
| By | |
| Distribution/ | |
| Availability Codes | |
| Dist | Avail and/or Special |
| A-1 | |

TIME DEPENDENCE OF HEAVY ION CONCENTRATION IN THE RING CURRENT

The focus of this project has been a comprehensive study of processes responsible for the decay of the storm time ring current. We have examined the importance to ring current ion loss of interaction with waves in the Pc 1 (Hz) and Pc 5 (mHz) frequency ranges, since such waves are a common feature of the ring current region during recovery from magnetic storms.

Pc 1 waves, identified as electromagnetic ion cyclotron (EMIC) waves, are generated by the loss cone distribution of ring current ions, analogous to their generation by radiation belt ions, which has been shown to determine the limit on stably trapped radiation belt ion fluxes (Kennel and Petchek, 1966). We have performed both linear dispersion analysis and particle simulations which show the importance of the loss cone vs. bi-Maxwellian nature of ring current ion distributions in determining the growth rate, angle of propagation and polarization of EMIC waves. In contrast to the conventional picture that EMIC waves are predominantly parallel propagating relative to B_0 , left hand polarized transverse ($\delta E \perp k$) modes, recent observations have

found that EMIC waves are commonly linearly polarized (Anderson et al., 1990). We have obtained a simple explanation for these observations, showing that a loss cone distribution generates obliquely propagating EMIC waves which are quasi-electrostatic, linearly polarized and analogous for ions to whistler modes on the electrostatic resonance cone (Denton et al., 1991).

We have shown that EMIC waves are effective at causing pitch angle diffusion of ring current H^+ into the loss cone, thereby providing a loss mechanism for enhanced storm time fluxes (Qian et al., 1990). EMIC waves are less effective at causing pitch angle diffusion of ring current O^+ , since O^+ is a minor constituent of the cold background plasma, and the O^+ cyclotron mode is not easily excited. This finding led us to the investigation of lower frequency modes in the Pc 5 range.

Pc 5 micropulsations are a completely different class of modes from EMIC waves. The latter are generated locally by the ring current ion loss cone, which is enhanced by global effects like drift shell splitting. The former is a global oscillation of the earth's magnetic field which may be driven internally by pressure gradients and particle resonances, or externally by changes in solar wind pressure at the magnetopause. We have concentrated on a class of internally driven, so called compressional Pc 5s which are commonly observed during the recovery phase of magnetic storms. Their frequency (2-5 mHz) is comparable to the bounce frequency of ring current O^+ and it has

been suggested that they may contribute to O^+ loss via a bounce resonance (Cladis and Kennel, 1986).

We have performed ring current test particle simulations in prescribed wave and background fields, including a dipole magnetic, convection and corotation electric field. We find that a drift-bounce resonant interaction causes ring current O^+ loss to the magnetopause in the energy range of a few tens of keV for moderate K_p (1.5 - 2). The compressional Pc 5s are standing waves along B_0 which propagate azimuthally in the direction of ring current ion drift. Thus it is possible for ring current ions in the right energy range to remain in phase with a single mode Pc 5 wave and either gain or lose energy from the wave fields. An ion which loses energy may not be able to overcome the convection potential across the dayside, as Figure 1 indicates. It will then $E \times B$ drift to the magnetopause.

While Pc 5s are often highly monochromatic, appearing almost sinusoidal in satellite magnetometer data, some spread in azimuthal mode number (perpendicular wavelength) is expected. If we assume a 10% spread in perpendicular wavelength, but the same total power in $\delta B/B_0$, the ion motion becomes diffuse in energy and L . The type of diffusion seen as a function of local time is apparent in the plots of ion number vs. energy in 2 keV bins shown in Fig. 2. One sees a bulk shift to lower energy due to the convection electric field as ions drift across the dayside, along with a spread to higher and lower energies associated with radial

diffusion. We have integrated the radial diffusion equation using a coefficient appropriate for the Pc 5 wave interaction described in Quarterly Report Number 9, and find good agreement with the numerical results (Li et al., 1991). Papers by Denton et al. and Li et al. are available upon request, and will be appended to Quarterly Report Number 17.





Figure 1



NPART=132 NOUT= 68 NPASS= 64

phim=95 f=0.002 bd=0.04 rkz=2.7 n=0 wphi0=0, r10=7, a0=2 mod=10
 econ=1 ecor=1 rkp=1.5 phi0=90 gphase0=0 gphased=0 am=16
 eki0=26, ekid=0.5 npe=11 pitch0=15 pitchd=20 nppitch=4
 rlp=6.0 rld=0.5 npl=3 numax=40000 err=1.e-05

Figure 2a







Figure 2b

References

- Anderson, B.J., K. Takahashi, R.E. Erlandson and L.J. Zanetti, Pc 1 pulsations observed by AMPTE/CCE in the earth's outer magnetosphere, Geophys. Res. Lett., 17, 1853, 1990.
- Cladis, J.B. and O.W. Lannartsson, On the loss of O^+ ions ($< 17\text{keV/e}$) in the ring current during the recovery phase of a storm, in Ion Acceleration in the Magnetosphere and Ionosphere, Tom Chang, ed. AGU, Washington, D.C., pp. 153-157, 1986.
- Denton, R.C., M.K. Hudson and I. Roth, Magnetospheric ion cyclotron waves with oblique propagation, submitted to J. Geophys. Res., 1991.
- Kennel, C.F. and H.E. Petchek, Limit on stably trapped particle fluxes, J. Geophys. Res., 71, 1, 1966.
- Li, X., A. Chan, M.K. Hudson and I. Roth, Loss of O^+ ions in the ring current due to interaction with Pc 5 waves, submitted to J. Geophys. Res., 1991.
- Qian, S., M.K. Hudson and I. Roth, Particle simulation of ion heating in the ring current, J. Geophys. Res., 95, 1001, 1990.